

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

2

## REPORT DOCUMENTATION PAGE

1a REF Ur		1b RESTRICTIVE MARKINGS	
2a SEC		3 DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release. Distribution unlimited.	
2b DEC		AD-A222 354	
4 PERFORMING ORGANIZATION REPORT NUMBER(S) 31		5 MONITORING ORGANIZATION REPORT NUMBER(S)	
6a NAME OF PERFORMING ORGANIZATION Massachusetts Institute of Technology	6b OFFICE SYMBOL (if applicable)	7a NAME OF MONITORING ORGANIZATION ONR	
6c ADDRESS (City, State, and ZIP Code) Department of Chemistry 77 Massachusetts Avenue Cambridge, MA 02139		7b ADDRESS (City, State, and ZIP Code) Office of the Navy Arlington, VA 22217	
8a NAME OF FUNDING/SPONSORING ORGANIZATION ONR	8b OFFICE SYMBOL (if applicable)	9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER	
8c ADDRESS (City, State, and ZIP Code) Office of the Navy Arlington, VA 22217		10 SOURCE OF FUNDING NUMBERS	
		PROGRAM ELEMENT NO. N-00014-32	TASK NO. K-0322
		WORK UNIT ACCESSION NO. P00-007	
11 TITLE (Include Security Classification) 1,2-DIMETHYL-1,2-DISILA-CLOSO-DODECABORANE(12), THE FIRST SILICON ANALOG OF AN ORTHO-CARBORANE			
12 PERSONAL AUTHOR(S) Dietmar Seyferth, Klaus Büchner, William S. Rees, Jr. and William M. Davis			
13a TYPE OF REPORT preprint	13b TIME COVERED FROM TO	14. DATE OF REPORT (Year, Month, Day) 1990-5-4	15. PAGE COUNT
16 SUPPLEMENTARY NOTATION communication in "Angewandte Chemie" & "Angewandte Chemie International Edition in English"			
17 COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	carboranes	
		silaboranes	
		organosilicon compounds	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) The reaction of $\text{CH}_3(\text{H})\text{Si}[\text{N}(\text{CH}_3)_2]_2$ with $\text{B}_{10}\text{H}_{14}$ in refluxing benzene gives 6,9- $[(\text{CH}_3)_2\text{NH}]_2\text{B}_{10}\text{H}_{12}$ and 1,2- $(\text{CH}_3)_2\text{Si}_2\text{B}_{10}\text{H}_{10}$ , <u>1</u> , in 15% yield. An X-ray diffraction study of the latter showed it has the ortho-carborane structure in which the two carbon atoms have been replaced by two silicon atoms. NMR spectroscopic ( $^{11}\text{B}$ , $^1\text{H}$ , $^{13}\text{C}$ , $^{29}\text{Si}$ ) and high resolution MS data are reported for <u>1</u> .			
20 DISTRIBUTION/AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION unclassified	
22a NAME OF RESPONSIBLE INDIVIDUAL Dr. S. Fishman		22b. TELEPHONE (Include Area Code) 202/696-4401	22c. OFFICE SYMBOL

DTIC  
ELECTE  
MAY 24 1990  
S E D

1,2-Dimethyl-1,2-disila-closo-dodecaborane(12), the First Silicon  
Analog of an ortho-Carborane\* \*

Dietmar Seyferth\*, Klaus Büchner, William S. Rees, Jr., William M. Davis

\* Prof. Dr. D. Seyferth, Dr. K. Büchner, Dr. W.S. Rees, Jr., Dr. W.M. Davis  
Department of Chemistry, Massachusetts Institute of Technology  
Cambridge, Massachusetts 02139 (USA)

\*\* This work was supported in part by the U.S. Office of Naval Research and  
by the award of a Feodor Lynen Fellowship to K.B. by the Alexander von  
Humboldt Foundation.

The icosahedral carboranes, especially those in which the two cluster  
carbon atoms are vicinal (*o*-carboranes), have received much attention since  
the first reports of their existence in 1963.<sup>(1)</sup> Notably absent in the many  
publications on this class of boron cluster compounds over the intervening 25  
odd years have been examples of icosahedral clusters in which the two mono-  
substituted carbon atoms of an *o*-carborane have been replaced by two mono-  
substituted silicon, germanium, tin or lead atoms. We report here the  
synthesis and structural characterization of 1, the first such Group IVA analog  
of an *o*-carborane: 1,2-dimethyl-1,2-disila-closo-dodecaborane(12) (or 1,2-  
dimethyl-*o*-silaborane). This compound is notable also in that it is the first  
example of a formal derivative of a disilyne,  $\text{CH}_3\text{Si}\equiv\text{SiCH}_3$  in the present case,

p 1

(Triple bonds)


Cont in  
the  
next page

a class of unsaturated organosilicon compounds of which there are to date no known examples. Keywords: Carboranes, Silaboranes, Organosilicon Compounds, organic chemistry, Silicon, Boron

The synthesis of 1,2-dimethyl-o-silaborane, 1, was the result of a deviation from the expected reaction course when a benzene solution of decaborane(14) and methylbis(dimethylamino)silane,  $\text{CH}_3(\text{H})\text{Si}[\text{N}(\text{CH}_3)_2]_2$ , 2, was heated at reflux under nitrogen. In prior work<sup>[2]</sup>, such a reaction of  $\text{B}_{10}\text{H}_{14}$  with  $(\text{CH}_3)_2\text{Si}[\text{N}(\text{CH}_3)_2]_2$  in refluxing benzene had resulted in formation of a linear polymer,  $[6,9-(\text{CH}_3)_2\text{NSi}(\text{CH}_3)_2\text{N}(\text{CH}_3)_2 \cdot \text{B}_{10}\text{H}_{12}]_x$ , a pale yellow, resinous material from whose melt fibers could be drawn and whose pyrolysis in a stream of ammonia to 1000°C gave boron nitride that contained 2-4% Si in high yield. However, the reaction of  $\text{B}_{10}\text{H}_{14}$  with 2 in refluxing benzene or toluene resulted in formation of a mixture of two white solids, 1 in 15% yield, based on  $\text{B}_{10}\text{H}_{14}$ , and 6,9- $[(\text{CH}_3)_2\text{NH}]_2\text{B}_{10}\text{H}_{12}$ , 3, in 58% yield (eq. 1). The former could be purified by sublimation at 90°C (0.01 Torr). Its yield has not been optimized. Compound 1 may be handled without exclusion of the laboratory atmosphere since it is stable to air and moisture.



Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



Very slow crystallization of 1 from dilute benzene solution at 3°C gave the 1:1 solvate, 1,2-(CH<sub>3</sub>)<sub>2</sub>Si<sub>2</sub>B<sub>10</sub>H<sub>10</sub>•C<sub>6</sub>H<sub>6</sub>, as rods whose structure was determined by single crystal X-ray diffraction.<sup>[3]</sup> The o-silaborane, like the o-carboranes, has a slightly distorted icosahedral structure (Fig. 1). The Si-Si bond distance of 2.308(2) Å is slightly less than that of "normal" Si-Si single bonds (~ 2.33 - 2.34 Å)<sup>[4]</sup>, but significantly longer than known Si-Si double bonds.<sup>[5]</sup> In comparison, the C-C bond distances in o-carboranes (1.655 Å, the mean value found in X-ray and gas phase electron diffraction studies<sup>[6]</sup>) are longer than the normal C(sp<sup>3</sup>)-C(sp<sup>3</sup>) distance of 1.54 Å. This is understandable in terms of the larger size of a silicon atom (covalent radius 1.18 Å), compared with a carbon atom (covalent radius 0.77 Å). The Si-B bond distances in 1 [2.017(3), 2.018(3), 2.113(4) and 2.116(3) Å] are very close to the sum of the covalent radii of Si and B, 2.07 Å (using 0.89 Å as the average boron

covalent radius in o-carboranes<sup>[6]</sup>). These distances may be compared to those of the Si-B bonds in *commo*-3,3'-Si(3,1,2-SiC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>)<sub>2</sub>, which contains a single silicon atom bonded within a 12-atom boron-containing cluster: 2.05(1) and 2.14(1) Å.<sup>[7]</sup>

The <sup>1</sup>H, <sup>13</sup>C, <sup>29</sup>Si and <sup>11</sup>B NMR spectra of 1 were recorded.<sup>[8]</sup> The <sup>11</sup>B NMR spectrum (160.367 MHz, acetone-d<sub>6</sub>, 25°C, external BF<sub>3</sub>•OEt<sub>2</sub>) was in agreement with the 1,2-disila-closo-dodecaborane(12) cage structure, showing four doublets in 2:2:2:4 intensity ration at δ<sub>B</sub> -11.2 (d, J = 147 Hz, B9, 12), -12.6 (d, J = 144 Hz, B8, 10), -13.5 (d, J = 152 Hz, B3, 6) and -14.5 (d, J = 147 Hz, B4, 5, 7, 11). The assignments are based on a 2D proton-decoupled <sup>11</sup>B-<sup>11</sup>B FT NMR spectrum. The CH<sub>3</sub> resonance was observed at -0.07 ppm in the <sup>1</sup>H NMR spectrum of 1 in C<sub>6</sub>D<sub>6</sub> solution. The proton-decoupled <sup>29</sup>Si NMR spectrum of 1 in C<sub>6</sub>D<sub>6</sub> solution at 50°C showed a singlet at -37.77 ppm, the proton-decoupled <sup>13</sup>C NMR spectrum, also in C<sub>6</sub>D<sub>6</sub> at 40°C, a singlet at -12.80 ppm. The upfield shifts observed in the <sup>13</sup>C and <sup>29</sup>Si spectra of 1 taken in C<sub>6</sub>D<sub>6</sub> solution, as compared with TMS, are consistent with the well-documented electron-deficient nature of boron cages.

Compound 1 is less stable than the o-carboranes. It melts at 201-203°C (sealed capillary) and decomposes with gas evolution above 230°C. It remains undecomposed when heated at reflux in wet tetrahydrofuran solution for several days. It was unaffected when treated with an excess of CF<sub>3</sub>CO<sub>2</sub>H in THF at room temperature and remained largely undecomposed when this solution was heated at reflux for 14 hr. It was found to be stable toward AlCl<sub>3</sub> in refluxing benzene. It reacts with and is degraded by alcoholic KOH in THF at room temperature within minutes and it decolorizes a solution of bromine in carbon tetrachloride after several hours at reflux.

Now that the first member of this new class of boron cluster compounds has been isolated, it may be expected that other 1,2-R<sub>2</sub>Si<sub>2</sub>B<sub>10</sub>H<sub>10</sub> compounds can be prepared. The development of their chemical reactivity at silicon and at boron will be of some interest as will a comparison of their chemistry with that of the o-carboranes. Our further research will deal with these questions.

## Experimental Procedure

A 250 ml three-necked flask equipped with a reflux condenser, an addition funnel and a magnetic stir-bar was flushed with argon and charged with 6.6 g (54.0 mmol) of  $B_{10}H_{14}$  and 120 ml of dry toluene. To the resulting solution was added dropwise at room temperature 7.1 g (53.7 mmol) of 2<sup>[9]</sup>. The reaction mixture was stirred and heated at reflux under argon for 24 hr. During this time a white solid precipitated. The reaction mixture was filtered and the filtrate evaporated at reduced pressure. The residue was extracted with two 100 ml portions of hot benzene. Filtration of the extracts and cooling to 3°C resulted in deposition of 1.45 g of 1 as an amorphous white powder that contained a small amount of white needles. Evaporation of the remaining benzene solution gave 0.6 g of 3, identified on the basis of its  $^1H$  and  $^{11}B$  NMR and mass spectra. The solid which had precipitated during the reaction was sublimed at 90°C (0.01 Torr) to give additional 0.15 g of 1. The sublimation residue was 6.0 g of 3. The total yield of the latter was 6.6 g, (58%). The total yield of 1 was 1.60g, a yield of 15% based on  $B_{10}H_{14}$ . Anal. Calcd. for  $C_2H_{16}Si_2B_{10}$ : C, 11.75; H, 7.89; Si, 27.48. Found: C, 11.64; H, 7.75; Si, 27.62. The high resolution MS showed molecular ion peaks at  $m/z$  206.1720

( $^{12}\text{C}_2\text{H}_{16}^{11}\text{B}_{10}^{28}\text{Si}_2$  calcd. 206.17211), 207.1715 ( $^{12}\text{C}_2\text{H}_{16}^{11}\text{B}_{10}^{28}\text{Si}^{29}\text{Si}$  calcd.

207.17168), 208.1688 ( $^{12}\text{C}_2\text{H}_{16}^{11}\text{B}_{10}^{28}\text{Si}^{30}\text{Si}$  calcd. 208.16898), and 209.1685

( $^{12}\text{C}_2\text{H}_{16}^{11}\text{B}_{10}^{29}\text{Si}^{30}\text{Si}$  calcd 209.16855).



1. Gmelin Handbook of Inorganic Chemistry, 8th Edition, "Borverbindungen", 2 (1974), 6 (1975), 11 (1977), and 12 (1977), and chapters in the first and second supplements (1980-1983), Springer-Verlag, Berlin. See also: R.N. Grimes, "Carboranes", Academic Press, New York, 1970, Chapter 6 and literature cited therein.
2. D. Seyferth and W.S. Rees, Jr., unpublished work.
3. The  $1,2-(\text{CH}_3)_2\text{Si}_2\text{B}_{10}\text{H}_{10} \cdot \text{C}_6\text{H}_6$  molecule crystallizes with crystallographic two-fold symmetry. One molecule of benzene was found for every molecule of 1 in two sites of equal occupancy; each site has the same crystallographic two-fold symmetry. Crystal data for 1  $\cdot \text{C}_6\text{H}_6$ : orthorhombic, space group Pccn (#56);  $a = 10.081(1)$ ,  $b = 10.666(8)$ ,  $c = 16.130(5)$  Å,  $V = 1734$  Å<sup>3</sup>,  $Z = 4$ ;  $\rho_{\text{calcd}} = 1.052$  g cm<sup>-3</sup>,  $F(000) = 592$ ,  $\mu(\text{MoK}\alpha) = 1.78$  cm<sup>-1</sup>, 2291 reflections measured, 2291 unique reflections, 1050 observed ( $I > 3\sigma(I)$ ), 89 variables,  $R = 0.044$ ,  $R_w = 0.058$ , Rigaku AFC6R diffractometer,  $\text{MoK}\alpha$  ( $\lambda = 0.71069$  Å). Further details of the crystal structure determination may be obtained from the Fachinformationszentrum Karlsruhe Gesellschaft für wissenschaftlich-technische Information mbH, D-7514 Eggenstein-Leopoldshafen 2 (FRG)


on quoting the deposition number \*\*\*\*\*, the names of the authors and the journal citation.

4. Representative Si-Si single bond distances: 2.340(9) Å in (CH<sub>3</sub>)<sub>6</sub>Si<sub>2</sub>: B. Beagley, J.J. Monaghan, T.G. Hewis, J. Mol. Struct. **8** (1971) 401; 2.338(4) Å in cyclo-[(CH<sub>3</sub>)<sub>2</sub>Si]<sub>6</sub>: H.L. Carrell, J. Donohue, Acta Cryst. **B28** (1972) 1566.
5. Si-Si double bond distances: 2.143(1) and 2.160(1) Å: M.J. Fink, M.J. Michalczyk, K.J. Haller, J. Michl, R. West, Organometallics **3** (1984) 793; 2.140(9) Å: S. Masamune, S. Murakami, J.T. Snow, H. Tobita, D.J. Williams, Organometallics **3** (1984) 3330.
6. V.S. Mastryukov, L.V. Vilkov, O.V. Dorofeeva, J. Mol. Struct. **24** (1975) 217.
7. W.S. Rees, Jr., D.M. Schubert, C.B. Knobler, M.F. Hawthorne, J. Am. Chem. Soc. **108** (1986) 5369.
8. NMR spectra, conditions: All measurements at 25 °C unless otherwise specified. <sup>11</sup>B, 160.37 and 96.235 MHz, BF<sub>3</sub>•OEt<sub>2</sub>; <sup>1</sup>H, 300 MHz, TMS; <sup>29</sup>Si, 99.34 MHz, TMS; <sup>13</sup>C, 75.43 MHz, TMS.
9. S.S. Washburne, W.R. Peterson, Jr., J. Organomet. Chem. **21** (1970) 59.

# CAPTION FOR FIGURE 1

ORTEP representation of 1,2-dimethyl-1,2-disila-*clos*-dodecaborane(12), drawn with 35% probability ellipsoids. Hydrogen atoms have been given arbitrary thermal parameters for clarity. Due to molecular symmetry, distances involving Si 2 are equivalent to those involving Si 1. Other symmetry related atoms are: C 1 and C 2; B 3 and B 6; B 4 and B 11; B 5 and B 7; B 8 and B 10; B 9 and B 12. Selected interatomic distances (in Å, with standard deviations of the last digit given in parentheses) are:

Si 1 - C 1	1.827(3)	B 3 - B 4	1.848(5)
Si 1 - Si 2	2.308(2)	B 4 - B 5	1.859(5)
Si 1 - B 3	2.116(3)	B 4 - B 8	1.774(5)
Si 1 - B 4	2.018(3)	B 4 - B 9	1.765(5)
Si 1 - B 5	2.017(3)	B 5 - B 6	1.849(5)
Si 1 - B 6	2.113(4)	B 5 - B 9	1.769(5)
B 6 - B 10	1.771(4)	B 5 - B 10	1.765(5)
B 8 - B 12	1.782(5)	B 9 - B 12	1.775(8)
B 10 - B 12	1.784(5)		



## ABSTRACT

The reaction of  $\text{CH}_3(\text{H})\text{Si}[\text{N}(\text{CH}_3)_2]_2$  with  $\text{B}_{10}\text{H}_{14}$  in refluxing benzene gives 6,9- $[(\text{CH}_3)_2\text{NH}]_2\text{B}_{10}\text{H}_{12}$  and 1,2- $(\text{CH}_3)_2\text{Si}_2\text{B}_{10}\text{H}_{10}$ , 1, in 15% yield. An X-ray diffraction study of the latter showed it has the ortho-carborborane structure in which the two carbon atoms have been replaced by two silicon atoms. NMR spectroscopic ( $^{11}\text{B}$ ,  $^1\text{H}$ ,  $^{13}\text{C}$ ,  $^{29}\text{Si}$ ) and high resolution MS data are reported for 1.